

# Early Termination of Convex QP Solvers in Mixed-Integer Programming for Real-Time Decision Making

Liang, Jiaming; Di Cairano, Stefano; Quirynen, Rien

TR2020-152 November 26, 2020

## Abstract

The branch-and-bound optimization algorithm for mixed-integer model predictive control (MI-MPC) solves several convex quadratic program relaxations, but often the solutions are discarded based on already known integer feasible solutions. This paper presents a projection and early termination strategy for infeasible interior point methods to reduce the computational effort of finding a globally optimal solution for MI-MPC. The method is shown to be also effective for infeasibility detection of the convex relaxations. We present numerical simulation results with a reduction of the total number of solver iterations by 42% for an MI-MPC example of decision making for automated driving with obstacle avoidance constraints.

*L-CSS with ACC 2021 Option*

© 2020 MERL. This work may not be copied or reproduced in whole or in part for any commercial purpose. Permission to copy in whole or in part without payment of fee is granted for nonprofit educational and research purposes provided that all such whole or partial copies include the following: a notice that such copying is by permission of Mitsubishi Electric Research Laboratories, Inc.; an acknowledgment of the authors and individual contributions to the work; and all applicable portions of the copyright notice. Copying, reproduction, or republishing for any other purpose shall require a license with payment of fee to Mitsubishi Electric Research Laboratories, Inc. All rights reserved.



# Early Termination of Convex QP Solvers in Mixed-Integer Programming for Real-Time Decision Making

Jiaming Liang<sup>1</sup>, Stefano Di Cairano<sup>1</sup> and Rien Quirynen<sup>1</sup>

**Abstract**—The branch-and-bound optimization algorithm for mixed-integer model predictive control (MI-MPC) solves several convex quadratic program relaxations, but often the solutions are discarded based on already known integer feasible solutions. This paper presents a projection and early termination strategy for infeasible interior point methods to reduce the computational effort of finding a globally optimal solution for MI-MPC. The method is shown to be also effective for infeasibility detection of the convex relaxations. We present numerical simulation results with a reduction of the total number of solver iterations by 42% for an MI-MPC example of decision making for automated driving with obstacle avoidance constraints.

**Index Terms**—Cyber-physical systems, mixed-integer programming, optimization algorithms.

## I. INTRODUCTION

MODEL predictive control (MPC) is a model-based control approach which allows to handle performance optimization and constraints by design [1]. This framework can be further extended to hybrid dynamical systems [2] that include both continuous and discrete decision variables, providing a powerful control design for a large class of problems, e.g., switched dynamical systems [3], discrete or quantized actuation [4], logic rules and temporal logic specifications [5]. The resulting optimization problems are non-convex because they contain variables that only take integer values. For a linear-quadratic objective, (piecewise) linear dynamics and linear constraints, the optimization problem can be formulated as a mixed-integer quadratic program (MIQP).

In this work, we aim at solving MIQPs of the form:

$$\min_{X, U} \sum_{k=0}^N \frac{1}{2} \begin{bmatrix} x_k \\ u_k \end{bmatrix}^\top H_k \begin{bmatrix} x_k \\ u_k \end{bmatrix} + \begin{bmatrix} x_k \\ u_k \end{bmatrix}^\top \begin{bmatrix} q_k \\ r_k \end{bmatrix} \quad (1a)$$

$$\text{s.t. } x_0 = \hat{x}_0, \quad (1b)$$

$$x_{k+1} = \begin{bmatrix} A_k & B_k \end{bmatrix} \begin{bmatrix} x_k \\ u_k \end{bmatrix} + a_k, \quad k \in \mathbb{Z}_0^{N-1}, \quad (1c)$$

$$y_k \leq \begin{bmatrix} C_k & D_k \end{bmatrix} \begin{bmatrix} x_k \\ u_k \end{bmatrix} \leq \bar{y}_k, \quad k \in \mathbb{Z}_0^N, \quad (1d)$$

$$u_{k,j} \in \mathbb{Z}, \quad j \in \mathcal{I}_k, \quad k \in \mathbb{Z}_0^N, \quad (1e)$$

where the notation  $\mathbb{Z}_a^b$  denotes the range of integers  $a, a + 1, \dots, b$ , and the optimization variables are the state  $X = [x_0^\top, \dots, x_N^\top]^\top$  and control trajectory  $U = [u_0^\top, \dots, u_N^\top]^\top$ . For simplicity of notation, the set  $\mathcal{I}_k$  denotes the indices of the discrete control variables for each step  $k \in \mathbb{Z}_0^N$  of the

prediction horizon. The Hessian matrices in (1a) are assumed to be positive semi-definite  $H_k \succeq 0$ , but not necessarily positive definite. Unlike standard MPC [1], our MIQP (1) can include discrete and/or continuous control variables on the terminal step of the MPC horizon.

MPC for hybrid dynamical systems aims at solving the MIQP (1) at every sampling time instant. In order to solve MIQPs to global optimality, most algorithms are based on a variant of the branch-and-bound (B&B) method [6]. The B&B strategy has been combined with various methods for solving the relaxed convex quadratic programs (QPs) in mixed-integer MPC (MI-MPC), such as dual active-set [7], primal active-set [8], interior point method (IPM) [9], dual projected gradient [10], and the alternating direction method of multipliers (ADMM) [11]. An advantage of a dual QP solver is that a dual feasible starting point for the child problems can be computed from the dual solution of the parent. This allows for early termination of the relaxed problem whenever the dual objective becomes larger than the current upper bound in the B&B [7], [10], [12].

IPMs are appealing for real-time optimization because they generally exhibit better worst-case performance than active-set type methods [13]. An early termination strategy for a dual feasible IPM is described in [14, Section 5.1]. Instead, here we focus on an infeasible IPM [15] that does not need a starting point that is dual feasible, and is applied to solve the convex relaxations within the B&B method. In general, all iterates of an infeasible primal-dual IPM are infeasible, although its limit points are both feasible and optimal [15]. Infeasible IPMs are appealing in B&B [16] because they can be effectively warm started, see [17], [18].

This paper proposes a first early termination strategy for an infeasible IPM to solve each convex QP relaxation in the B&B method, without the assumption of strict positive definiteness of the Hessian in (1) and without the need to compute a dual feasible starting point. We developed a computationally efficient projection to compute a dual feasible solution guess for early termination before the optimum of the QP is found. We additionally show that our early termination strategy is effective for infeasibility detection for the convex relaxations in the B&B method.

The paper is organized as follows. Section II summarizes background materials on optimization and B&B methods. Section III presents our projection and early termination strategy for primal-dual IPMs, and the effectiveness for infeasibility detection is motivated in Section IV. Section V presents numerical results for an MI-MPC example of automated driving using the active-set based IPM (ASIPM) method with

<sup>1</sup>Mitsubishi Electric Research Laboratories, Cambridge, MA, 02139, USA. {liang, dicairano, quirynen}@merl.com

warm starting in [18]. The conclusions are summarized in Section VI.

## II. PROBLEM FORMULATION

We reformulate the MIQP in (1) in the compact notation

$$\min_z \frac{1}{2} z^\top H z + h^\top z \quad (2a)$$

$$\text{s.t. } C z \leq c, \quad F z = f, \quad (2b)$$

$$z_j \in \mathbb{Z}, \quad j \in \mathcal{I}, \quad (2c)$$

where  $z$  includes all primal optimization variables and the set  $\mathcal{I}$  indexes the integer variables. For many practical MPC formulations, the objective can directly be written as

$$\frac{1}{2} z^\top H z + h^\top z = \frac{1}{2} x^\top Q x + h_x^\top x + h_y^\top y, \quad (3)$$

where  $H \succeq 0$  and  $Q \succ 0$ , by partitioning  $z$  into  $x$  and  $y$ , entering in the linear-quadratic and linear-only terms, respectively. The objective of any MIQP (2) can always be reformulated as in (3) by a change of variables, e.g., based on the eigenvalue decomposition for the Hessian matrix  $H \succeq 0$ .

### A. Branch-and-bound Optimization

The standard B&B algorithm [6] sequentially creates partitions for the search space of the original mixed-integer program and compute bounds for those partitions. Each integer-feasible solution provides a global upper bound to the optimum. Local lower bounds can be computed from convex relaxations of the MIQP for a particular partition or node of the B&B tree. The B&B effectiveness stems from pruning certain partitions or nodes whenever the local lower bound exceeds the current global upper bound.

The B&B partitions are constructed through branching, and selecting an appropriate integer variable for branching is a key B&B step, e.g., studied extensively in [19]. Another ingredient is the pre-solve routine to construct tighter convex relaxations leading to less branching, see the overview in [20]. For lowering the upper bound, heuristics are often used to compute integer-feasible solutions, e.g., by a feasibility pump [21] or warm starting [22]. The B&B method used in this paper for numerical tests is similar to [8] and exploits domain propagation in the pre-solve routine, reliability branching based on pseudo-cost, and warm starting.

### B. Convex QP Relaxations

We focus on the efficient solution of convex QP relaxations in a B&B optimization method. For a particular node of the B&B tree, a convex QP is obtained by relaxing the integer equality constraint in (2c) as

$$z_j \leq z_j \leq \bar{z}_j, \quad j \in \tilde{\mathcal{I}}, \quad (4)$$

where the index set  $\tilde{\mathcal{I}}$  denotes each integer variable that has not been fixed due to branching, or due to the pre-solve routine, in the current node of the B&B tree. The values  $z_j$  and  $\bar{z}_j$  denote the lower and upper bound values for each integer variable  $j \in \tilde{\mathcal{I}}$ , respectively. As discussed in [20], fixed variables, i.e.,

$z_j = \bar{z}_j$ , and redundant constraints should be removed from each convex QP for efficiency.

We do not need to solve a convex subproblem if

- 1) the convex QP relaxation is infeasible,
- 2) the optimal solution is detected to have an objective value that exceeds the current global upper bound.

In both cases, the node, and hence the corresponding subtree, can be pruned from the B&B tree to find the optimal solution. A considerable computational effort can be avoided if the above scenarios are detected early, i.e., more quickly than the solving time of the convex QP relaxations. In the present paper, we propose a tailored early termination strategy for infeasible primal-dual IPMs to handle both cases and to reduce the computational effort of the B&B method without affecting the quality of the optimal solution.

### C. Dual QP Problem Formulation

We consider the primal convex QP of the form

$$\min_{x,y} \phi(x,y) := \frac{1}{2} x^\top Q x + h_x^\top x + h_y^\top y \quad (5a)$$

$$\text{s.t. } G_x x + G_y y \leq g, \quad (5b)$$

$$F_x x + F_y y = f, \quad (5c)$$

where  $Q \succ 0$  in the primal objective  $\phi(x,y)$ , and the inequality constraints (5b) include both the original inequalities from (2b) and the convex relaxations (4) of the integer constraints. We additionally use the compact notation  $z := [x^\top y^\top]^\top$ ,  $G := [G_x | G_y]$ ,  $F := [F_x | F_y]$ ,  $h := [h_x^\top h_y^\top]^\top$  and  $H := \begin{bmatrix} Q & 0 \\ 0 & 0 \end{bmatrix}$ . The dual QP of (5) reads as

$$\max_{\mu,\lambda} \psi(\mu,\lambda) := -\frac{1}{2} \|\hat{h}(\mu,\lambda)\|_{Q^{-1}}^2 - \begin{bmatrix} g \\ f \end{bmatrix}^\top \begin{bmatrix} \mu \\ \lambda \end{bmatrix} \quad (6a)$$

$$\text{s.t. } G_y^\top \mu + F_y^\top \lambda = -h_y, \quad (6b)$$

$$\mu \geq 0, \quad (6c)$$

where  $\lambda$  and  $\mu$  denote the Lagrange multipliers for the equality and inequality constraints, respectively, and

$$\hat{h}(\mu,\lambda) := h_x + G_x^\top \mu + F_x^\top \lambda, \quad (7)$$

for simplifying the dual objective function  $\psi(\mu,\lambda)$  in (6a).

### D. Primal-dual Interior Point Method

A primal-dual IPM uses a Newton-type method to solve a sequence of relaxed Karush-Kuhn-Tucker (KKT) conditions for the convex QP in (5). An iteration of the IPM usually solves the reduced linear system

$$\begin{bmatrix} H & F^\top & G^\top \\ F & 0 & 0 \\ G & 0 & -W^k \end{bmatrix} \begin{bmatrix} \Delta z^k \\ \Delta \lambda^k \\ \Delta \mu^k \end{bmatrix} = - \begin{bmatrix} r_z^k \\ r_\lambda^k \\ \bar{r}_\mu^k \end{bmatrix}, \quad (8)$$

where  $W^k = \text{diag}(w^k) \succ 0$  and  $w_i^k = s_i^k / \mu_i^k > 0$ . The right-hand side in (8) denotes the residual value for the optimality conditions and reads as

$$\begin{aligned} r_z^k &= H z^k + F^\top \lambda^k + G^\top \mu^k + h, \\ r_\lambda^k &= F z^k - f, \quad r_\mu^k = G z^k - g + s^k, \\ r_s^k &= M^k S^k \mathbf{1} - \tau^k \mathbf{1}, \quad \bar{r}_\mu^k = r_\mu^k - M^{k-1} r_s^k, \end{aligned} \quad (9)$$



projection step in (13) may be needed to ensure dual feasibility  $\|F_y^\top \lambda + G_y^\top \mu + h_y\| < tol$  when using the inexact Newton implementation of [18].

---

**Algorithm 1** Early termination for IPM in B&B method.

---

```

1: Input: Warm start  $\{(z^0, \mu^0, \lambda^0, s^0)\}$ ,  $tol$ , and  $UB$ .
2: while  $\max\{\tau^k, \|r^k\|\} > tol$  do
3:   if  $\psi(\mu^k, \lambda^k) > UB$  & dual_feasible then
4:     break while loop. ▷ Early termination
5:   else if  $\psi(\mu^k, \lambda^k) > UB$  then
6:     Compute projection step  $(\Delta\mu, \Delta\lambda)$  in (13).
7:      $\mu \leftarrow \mu^k + \Delta\mu$ ,  $\lambda \leftarrow \lambda^k + \Delta\lambda$ , and
8:      $r_y \leftarrow F_y^\top \lambda + G_y^\top \mu + h_y$ .
9:     if  $\mu > 0$  &  $\|r_y\| < tol$  then
10:       $\mu^k \leftarrow \mu$ ,  $\lambda^k \leftarrow \lambda$ ,  $r_y^k \leftarrow r_y$ , and
11:      dual_feasible  $\leftarrow 1$ .
12:      if  $\psi(\mu^k, \lambda^k) > UB$  then
13:        break while loop. ▷ Early termination
14:      end if
15:    end if
16:  end if
17:  Perform an IPM iteration (8), e.g., see [18].
18: end while

```

---

### C. Computational Complexity

The proposed early termination strategy requires two computational steps, i.e., the projection and the dual objective evaluation, which are typically not needed in a standard IPM. Considering the optimal control structured program (1), the evaluation of the dual objective value (6a) requires

$$N(n^2 + 2nm + 2np), \quad (17)$$

operations to compute  $L^{-1}(G_x^\top \mu + F_x^\top \lambda)$  based on a block-diagonal Cholesky factorization  $Q = LL^\top$ , in which  $n$ ,  $m$  and  $p$  are the number of variables  $x$ , the number of equality and inequality constraints per control interval, respectively.

Based on (13), we can perform one projection step at the computational cost of one IPM iteration. This allows the reuse of the corresponding matrix factorization in the subsequent IPM iteration if the projection is not successful. Based on the particular ASIPM implementation, as proposed recently in [18], one iteration requires a block-tridiagonal Cholesky factorization for the matrix in (14), for which the dominant terms in the computational cost are

$$N\left(\frac{7}{3}n_x^3 + 4n_x^2n_u + 2n_xn_u^2 + \frac{1}{3}n_u^3\right), \quad (18)$$

where  $n_x$  and  $n_u$  denote the number of state and control variables per interval in (1), respectively. Then, the linear system in (13) can be solved by

$$N(6n_x^2 + 8n_xn_u + 2n_u^2 + 2(n_x + n_u)(m + p)), \quad (19)$$

operations for the resulting block-structured forward and backward substitution. Since  $n \leq (n_x + n_u)$ , and often  $n \ll (n_x + n_u)$  due to many auxiliary variables in hybrid systems [2]

for which the Hessian contribution is zero, the cost for a dual objective evaluation (17) is considerably smaller than the projection cost (19), as anticipated.

## IV. PRIMAL INFEASIBILITY DETECTION

We first present a certificate of primal infeasibility and then justify why our proposed early termination strategy can be also effective for primal infeasibility detection. We confine ourselves to the compact form of the QP relaxation in Eq. (5) and a primal-dual IPM described in Section II-D.

### A. Certificate of Primal Infeasibility

Let the primal problem (5) with strict inequalities,

$$Gz < g, \quad Fz = f, \quad (20)$$

be infeasible. Using Farkas' lemma [15], we can derive the following certificate of strict infeasibility for (20),

$$G^\top \tilde{\mu} + F^\top \tilde{\lambda} = 0, \quad g^\top \tilde{\mu} + f^\top \tilde{\lambda} < 0, \quad \tilde{\mu} > 0. \quad (21)$$

Eq. (20) is strictly infeasible if and only if there exists a pair  $(\tilde{\mu}, \tilde{\lambda})$  satisfying (21). If  $(\hat{\mu}, \hat{\lambda})$  is a dual feasible solution satisfying (6b) and (6c), then  $(\mu, \lambda) = (\hat{\mu} + \alpha\tilde{\mu}, \hat{\lambda} + \alpha\tilde{\lambda})$  also satisfies (6b) and (6c) for  $\alpha > 0$ , and the dual objective in (6a)  $\rightarrow \infty$  as  $\alpha \rightarrow \infty$ , i.e., the dual problem is unbounded.

### B. Early Termination for Infeasible QPs

As discussed in detail in [24], the standard iterates of an IPM can be used to generate certificates of infeasibility (21). Intuitively, since the dual objective is unbounded in case of primal infeasibility, we can terminate the solver whenever the dual objective of a dual feasible solution is ‘‘large enough’’. We motivate why our proposed early termination strategy is effective for infeasibility detection and, given a tight UB from the B&B optimization method, it may lead to termination before a certificate of infeasibility is found.

Proposition 4.3 will prove that the dual objective  $\psi(\mu^k, \lambda^k)$  for the primal-dual IPM iterates is unbounded in the limit. Before, we need to prove two technical lemmas.

*Lemma 4.1:* The following statements about the IPM iterates (8) hold for  $k \geq 0$ :

a) for some  $\beta_k \in [0, 1]$ , iterates  $z^k$  and  $s^k$  satisfy

$$M \begin{bmatrix} z^k \\ s^k \end{bmatrix} = \beta_k M \begin{bmatrix} z^0 \\ s^0 \end{bmatrix} + (1 - \beta_k) \begin{bmatrix} g \\ f \end{bmatrix} =: r_{z,s}^k, \quad (22)$$

where  $M := \begin{bmatrix} G & I \\ F & 0 \end{bmatrix}$ , and  $r_{z,s}^k$  is bounded;

b)  $h^k := Hz^k + G^\top \mu^k + F^\top \lambda^k$  is bounded.

*Proof:* a) Eq. (22) holds for  $k = 0$  with  $\beta_0 = 1$ . Next, assuming that (22) holds for  $k \geq 0$ , it follows from Eq. (8) and (9), and  $\Delta s^k = -(W^k \Delta \mu^k + M^{k-1} r_s^k)$  that

$$M \begin{bmatrix} z^{k+1} \\ s^{k+1} \end{bmatrix} = (1 - \alpha^k) M \begin{bmatrix} z^k \\ s^k \end{bmatrix} + \alpha^k \begin{bmatrix} g \\ f \end{bmatrix}, \quad (23)$$

where  $(z^{k+1}, s^{k+1}) = (z^k + \alpha^k \Delta z^k, s^k + \alpha^k \Delta s^k)$ , and  $\alpha^k \in (0, 1]$  is the stepsize taken in the IPM. Hence, it follows from (23) that Eq. (22) holds for  $k+1$  with  $\beta_{k+1} = \beta_k(1 - \alpha^k)$ .

It follows from its definition in (22) that  $r_{z,s}^k$  is bounded as a convex combination of two bounded vectors.

b) This statement follows from similar arguments as for the derivations in the proof of (a). ■

It is easy to see from (22) that  $\beta_k > 0$  in case of strict primal infeasibility. The following lemma confirms this observation and the boundedness of  $s^k$  and  $z^k$ .

*Lemma 4.2:* If the primal problem (20) is strictly infeasible, then the following statements hold for  $k \geq 0$ :

a) there exists  $\bar{\mu} > 0$  such that

$$\beta_k \geq (1 + \bar{\mu}^\top s^0)^{-1} > 0, \quad \bar{\mu}^\top s^k \leq \bar{\mu}^\top s^0; \quad (24)$$

b) assuming that  $[G^\top F^\top]^\top$  has full rank, then the iterates  $s^k$  and  $z^k$  are bounded.

*Proof:* a) Certificate (21) is equivalent to the existence of  $\bar{\mu}$  and  $\bar{\lambda}$  satisfying

$$G^\top \bar{\mu} + F^\top \bar{\lambda} = 0, \quad g^\top \bar{\mu} + f^\top \bar{\lambda} = -1, \quad \bar{\mu} > 0. \quad (25)$$

Following a similar argument as in the proof [24, Prop. 5.1], i.e., multiplying (22) by  $[\bar{\mu}^\top \bar{\lambda}^\top]$  from left, and using (25) and the fact that  $s^k > 0$ , we have

$$\beta_k (1 + \bar{\mu}^\top s^0) - 1 = \bar{\mu}^\top s^k > 0, \quad (26)$$

and hence the first inequality in (24) holds. Moreover, the second inequality in (24) follows from (26) and  $0 < \beta_k \leq 1$ .

b) Using the second inequality in (24) and  $\bar{\mu} > 0$ ,  $s^k > 0$  and  $s^0 > 0$ , we know that  $s^k$  is bounded. From (22), using Lemma 4.1(a) and  $s^k$  being bounded,  $z^k$  is also bounded because  $[G^\top F^\top]^\top$  has full rank. ■

We can now prove the main result of this section, namely, if the primal problem is infeasible, the dual objective of the IPM tends to infinity, which allows for early termination.

*Proposition 4.3:* If the sequence of IPM iterates  $\{(z^k, \mu^k, \lambda^k, s^k)\}$  satisfy  $\mu^{k^\top} s^k \leq \mu^{0^\top} s^0$  and  $\|\mu^k\| \rightarrow \infty$ , then the dual objective  $\psi(\mu^k, \lambda^k) \rightarrow \infty$ .

*Proof:* Lemmas 4.1(b) and 4.2(b) imply that  $\hat{h}(\mu^k, \lambda^k)$  in (7) is bounded. As a result, in order to show that the dual objective  $\psi(\mu^k, \lambda^k)$  (see (6a)) is unbounded for the IPM iterates, it suffices to show

$$\ell^k := g^\top \mu^k + f^\top \lambda^k \rightarrow -\infty. \quad (27)$$

Multiplying (22) by  $[\mu^{k^\top} \lambda^{k^\top}]$  from the left, using the assumption that  $\mu^{k^\top} s^k \leq \mu^{0^\top} s^0$ , the definition of  $\ell^k$  in (27), and the identity  $G^\top \mu^k + F^\top \lambda^k = h^k - H z^k$  (see Lemma 4.1(b)), after rearranging the terms, we obtain

$$(1 - \beta_k) \ell^k \leq t^k + (\mu^0 - \beta_k \mu^k)^\top s^0, \quad (28)$$

where  $t^k := (z^k - \beta_k z^0)^\top (h^k - H z^k)$ . Lemmas 4.1(b) and 4.2(b) imply that  $t^k$  is bounded. For  $\|\mu^k\| \rightarrow \infty$ , since  $\mu^k > 0$  and  $s^0 > 0$ , we have  $\mu^{k^\top} s^0 \rightarrow \infty$ . It follows from  $\beta_k > 0$  in Lemma 4.2(a) that the right-hand side of (28) tends to  $-\infty$ . Finally, since  $\beta_k < 1$  in (22), inequality (28) implies (27) and concludes the proof. ■

## V. NUMERICAL EXAMPLE

Next, we provide numerical simulation results to illustrate the performance of our early termination strategy applied to the ASIPM QP solver with warm starting [18], as part of the B&B implementation from [8] in MATLAB.

### A. Problem Formulation: Vehicle Decision Making

The numerical example involves an MI-MPC formulation of a high-level motion planning task for an autonomous vehicle, including discrete decisions resulting from lane changes, static and dynamic obstacles. Similar to the problem formulation in [5], the vehicle motion is modeled in curvilinear coordinates resulting in affine system dynamics with 3 state variables and 2 control inputs. Additional decision variables are defined to model lane changes and obstacle avoidance, resulting in an MIQP formulation of the form in (1) with  $N = 15$  intervals,  $n_x = 6$  state variables and  $n_u = 18$  control variables, including 14 binary decision variables per control interval. Due to the pre-solve routine based on domain propagation [8], many of the 224 binary decision variables can be eliminated in practice.

### B. Numerical Simulation Results

The performance of the early termination strategy in Algorithm 1 for this MI-MPC example is presented in Fig. 1, including the total number of QP calls and solver iterations at each time step, as well as the reduction of computational effort in percentage. The x-axis of each subplot is the simulation time in  $[0, 170]$  s, using a sampling period of 2 s. It can be observed that the reduction in QP iterations is most considerable when a larger number of QPs need to be solved in the B&B method to compute the global solution of the MIQP. This is where it matters the most for real-time applications that must consider the worst-case execution time. Overall, in this particular numerical example, early termination reduces 42% of the total QP iterations.

Fig. 2 presents a detailed comparison between early terminated and fully solved QPs to solve the MIQP at each time step. Subplot 1 compares the number of early terminated and fully solved QPs, and shows that early termination of the IPM happens often. More specifically, about 36% of all QPs are early terminated in this particular example. Subplot 2 compares the average number of iterations in early terminated QPs with and without warm starting, and shows that early termination benefits from warm starting. In fact, the number of QP iterations is 0 at times 112 – 134 s, i.e., the warm start leads to immediate termination of the IPM. Subplot 3 presents the average number of iterations in the solved QPs with and without warm starting. Comparing subplots 2 and 3, one can observe that early terminated QPs take considerably less iterations than fully solved QPs.

Table I finally illustrates the performance of our early termination strategy for infeasibility detection of three infeasible QPs from the MI-MPC example. We compare the number of IPM iterations to obtain a certificate of infeasibility (21) versus the early termination strategy from Algorithm 1, with




Fig. 1. QP calls and total iterations with and without early termination: early termination reduces total number of QP iterations by 42%.




Fig. 2. Early terminated versus fully solved QPs: average number of QP iterations in both cases, with and without warm starting of ASIPM.

and without warm starting of ASIPM [18]. Table I shows that early termination requires considerably less IPM iterations than the computation of a certificate of infeasibility. Note that early termination does not distinguish between a case of primal infeasibility or a case where the objective value exceeds the current UB. In addition, Table I shows that warm starting can reduce the number of IPM iterations further and it can lead to immediate termination, i.e., without the need to perform any IPM iterations.

TABLE I

INFEASIBILITY DETECTION: # OF QP ITERATIONS FOR CERTIFICATE VERSUS EARLY TERMINATION WITH AND WITHOUT WARM STARTING.

	QP # 1	QP # 2	QP # 3
Certificate of primal infeasibility	40	45	38
Early termination: cold started	10	12	10
Early termination: warm started	0	0	11

## VI. CONCLUSIONS

We proposed an efficient early termination strategy based on a projection step tailored to IPMs, in order to reduce the computational cost within the B&B method in solving MI-MPC problems. In addition, we proved that our early termination strategy can be effective for infeasibility detection and

often it terminates the IPM before a certificate of infeasibility is found. Finally, we illustrated the performance of this early termination technique based on numerical simulation results for an MI-MPC example of vehicle decision making.

## REFERENCES

- [1] D. Mayne and J. Rawlings, *Model Predictive Control*. Nob Hill, 2013.
- [2] A. Bemporad and M. Morari, “Control of systems integrating logic, dynamics, and constraints,” *Automatica*, vol. 35, pp. 407–427, 1999.
- [3] T. Marcucci and R. Tedrake, “Mixed-integer formulations for optimal control of piecewise-affine systems,” in *Proc. of the 22nd ACM Inter. Conf. on Hybrid Systems*, New York, NY, USA, 2019, p. 230–239.
- [4] A. Walsh, S. Di Cairano, and A. Weiss, “MPC for coupled station keeping, attitude control, and momentum management of low-thrust geostationary satellites,” in *Proc. American Control Conf. (ACC)*, 2016, pp. 7408–7413.
- [5] Y. E. Sahin, R. Quirynen, and S. Di Cairano, “Autonomous vehicle decision-making and monitoring based on signal temporal logic and mixed-integer programming,” in *Proc. American Control Conf. (ACC)*, 2020, pp. 454–459.
- [6] C. A. Floudas, *Nonlinear and mixed-integer optimization: fundamentals and applications*. Oxford University Press, 1995.
- [7] D. Axehill and A. Hansson, “A mixed integer dual quadratic programming algorithm tailored for MPC,” in *Proc. IEEE Conf. Decision and Control (CDC)*. IEEE, 2006, pp. 5693–5698.
- [8] P. Hespanhol, R. Quirynen, and S. Di Cairano, “A structure exploiting branch-and-bound algorithm for mixed-integer model predictive control,” in *Proc. European Control Conf. (ECC)*, 2019, pp. 2763–2768.
- [9] D. Frick, A. Domahidi, and M. Morari, “Embedded optimization for mixed logical dynamical systems,” *Computers & Chemical Engineering*, vol. 72, pp. 21–33, 2015.
- [10] V. V. Naik and A. Bemporad, “Embedded mixed-integer quadratic optimization using accelerated dual gradient projection,” *IFAC-PapersOnLine*, vol. 50, no. 1, pp. 10723–10728, 2017.
- [11] B. Stellato, V. Naik, A. Bemporad, and P. Goulart, “Embedded mixed-integer quadratic optimization using the OSQP solver,” in *Proc. European Control Conf. (ECC)*, 2018.
- [12] R. Fletcher and S. Leyffer, “Numerical experience with lower bounds for MIQP branch-and-bound,” *SIAM Journal on Optimization*, vol. 8, no. 2, pp. 604–616, 1998.
- [13] R. Bartlett, A. Wächter, and L. Biegler, “Active Set vs. Interior Point Strategies for Model Predictive Control,” in *Proc. American Control Conf. (ACC)*, Chicago, IL, 2000, pp. 4229–4233.
- [14] J. E. Mitchell, P. M. Pardalos, and M. G. C. Resende, “Interior point methods for combinatorial optimization,” in *Handbook of Combinatorial Optimization*. Boston, MA: Springer US, 1998, pp. 189–297.
- [15] S. J. Wright, *Primal-Dual Interior-Point Methods*. Philadelphia: SIAM Publications, 1997.
- [16] P. Munari and J. Gondzio, “Using the primal-dual interior point algorithm within the branch-price-and-cut method,” *Computers & Operations Research*, vol. 40, no. 8, pp. 2026 – 2036, 2013.
- [17] A. Engau, M. F. Anjos, and A. Vannelli, “On interior-point warmstarts for linear and combinatorial optimization,” *SIAM Journal on Optimization*, vol. 20, no. 4, pp. 1828–1861, 2010.
- [18] J. Frey, S. Di Cairano, and R. Quirynen, “Active-set based inexact interior point QP solver for model predictive control,” in *Proc. IFAC World Congr.*, 2020.
- [19] T. Achterberg, T. Koch, and A. Martin, “Branching rules revisited,” *Operations Research Letters*, vol. 33, no. 1, pp. 42–54, 2005.
- [20] T. Achterberg, R. E. Bixby, Z. Gu, E. Rothberg, and D. Weninger, “Presolve reductions in mixed integer programming,” *ZIB Report*, pp. 16–44, 2016.
- [21] T. Achterberg and T. Berthold, “Improving the feasibility pump,” *Discrete Optimization*, vol. 4, no. 1, pp. 77–86, 2007.
- [22] A. Bemporad and V. V. Naik, “A numerically robust mixed-integer quadratic programming solver for embedded hybrid model predictive control,” in *Proc. 6th IFAC NMPC Conf.*, Madison, USA, 2018.
- [23] S. Boyd and L. Vandenberghe, *Convex Optimization*. Cambridge: University Press, 2004.
- [24] M. J. Todd, “Detecting infeasibility in infeasible-interior-point methods for optimization,” in *Foundations of computational mathematics*. Cambridge Univ. Press, 2004, pp. 157–192.